

## Construction of a BAC library for *Haplochromis chilotes*, a cichlid fish from Lake Victoria

Masakatsu Watanabe<sup>1</sup>, Naoki Kobayashi<sup>1</sup>, Asao Fujiyama<sup>2,3</sup>,  
and Norihiro Okada<sup>1,\*</sup>

<sup>1</sup>Graduate School of Bioscience and Biotechnology, Tokyo Institute of Technology, 4259,  
Nagatsuta-cho, Midori-ku, Yokohama, 226-8501, Japan

<sup>2</sup>Information Research Division, National Institute of Informatics, 2-1-2,  
Hitotsubashi, Chiyoda-ku, Tokyo, 101-8430, Japan

<sup>3</sup>RIKEN Genomic Sciences Center, 1-7-22 Suehiro-cho, Tsurumi-ku,  
Yokohama, 230-0045, Japan.

(Received 10 October 2002, accepted 9 January 2003)

Cichlid fishes in Lake Victoria are model organisms for studying rapid radiation and speciation. On the way to examine the molecular basis of how these cichlid fishes achieved such a remarkable morphological diversification, we constructed a bacterial artificial chromosome (BAC) library derived from a cichlid species, *Haplochromis chilotes*, from Lake Victoria. The library includes 157,056 clones with the average insert size of 128 kb, corresponding to a 10-fold coverage of the *H. chilotes* genome. Given that the cichlid fishes endemic to Lake Victoria are closely related to one another phylogenetically and their genomes are nearly identical, this BAC library can be utilized to isolate genes from the more than 200 Haplochromine cichlid species in Lake Victoria.

**Key words:** cichlid, BAC library, Lake Victoria

BAC libraries (Shizuya et al., 1992) are very useful for analyzing genome structure in almost all organisms. To date, many BAC libraries have been constructed, and some have been utilized for genome sequencing projects. Regarding a cichlid BAC library, Katagiri et al. (2001) constructed a series of four BAC libraries using Nile tilapia (*Oreochromis niloticus*). These BACs cover 2–64x of the genome, and have been utilized in the cichlid genome project. Nile tilapia is recognized as an important commercial food source and is presently cultivated in more than 65 countries. From an evolutionary viewpoint, the tilapia constitutes an ancient lineage of East African cichlids, but displays less morphological diversity than the Haplochromine cichlids (Fryer and Iles, 1972; Greenwood, 1984; Stiassny, 1991). In East Africa, there are three Great Lakes, namely Lakes Tanganyika, Malawi and Victoria within which reside >1000 morphologically diverse endemic cichlid species that have adapted to these ecosystems over 10 My of evolution. In particular, the Haplochromine cichlid species in Lake Victoria are regarded as model organisms of adaptive radiation and rapid speciation (Barel et al., 1991; Johnson et al., 1996).

We constructed a BAC library derived from *Haplochromis*

*chilotes* of Lake Victoria. Genomic DNA was prepared from 10<sup>9</sup> reticulocytes isolated from two *H. chilotes* specimens. After anesthetizing the fish with 3-aminobenzoic acid ethyl ester (MS-222, Sigma, USA), arterial blood was collected and mixed with an equal volume of buffer (10 mM Tris-HCl [pH 8.0], 0.5 M EDTA). The reticulocytes were washed twice with phosphate-buffered saline and then treated with lysis buffer (10 mM Tris-HCl pH 8.0, 50 mM NaCl, 1% lithium dodesyl sulfate, 100mM EDTA) (Amemiya et al., 1996) for 2 hr at 37°C in 1.0% agarose gel plugs, each plug containing 5 × 10<sup>7</sup> reticulocytes, then stored in 20% NDS solution consists of 0.2% N-laurylsarcosine, 2 mM Tris-HCl (pH 9.0), 0.14 M EDTA. After exchanging the NDS buffer with TE buffer (10 mM Tris-HCl [pH 8.0], 0.1 mM EDTA), the genomic DNA was then partially digested with *Sac* I and 150–250 kb DNA fragments were isolated using pulse field gel electrophoresis. The DNA fragments eluted from the gel were ligated into pKS145 (Fujiyama et al., 2002) and an aliquot of the ligation reaction was used to transform *E. coli* DH10B. Using Flexys robot (Genomic Solutions, USA) and 3D:Biomek FX robot (BECKMAN COULTER, USA), a total of 157,056 BAC clones were chosen and arrayed in 409 × 384-well microtiter plates in LB medium containing 10% glycerol and 25 µg/ml ampicillin. The plates were incubated overnight at 37°C and then stored at –80°C. The BAC library was constructed so as to be

Edited by Hidenori Tachida

\* Corresponding author. E-mail: nokada@bio.titech.ac.jp




Fig. 1. Analysis of the average size of the BAC library inserts. The insert sizes of 20 randomly selected BAC clones were analyzed by pulse field gel electrophoresis. After digestion with *Not* I, an aliquot of each digest was applied to the gel. Each lane contains an insert with a vector fragment (7.4 kb). Electrophoresis was performed by CHEF-DR III (Bio-Rad) in 0.5x TBE buffer at 160 V for 15 hr at 16°C.

amenable to the dimension pooling system for PCR screening. To estimate the size of the average DNA insert in the BAC library, 80 randomly selected clones were digested with *Not* I and analyzed by pulse field gel electrophoresis. Only 20 clones are shown in Fig. 1. The average insert size was 128 kb, corresponding to 10x coverage of the *H. chilotes* genome. The genome size is assumed by that of *Oreochromis niloticus* (Majumdar and McAndrew, 1986). As a pilot screening experiment, we obtained independent seven BAC clones of the *lws* (a long wavelength-sensitive) opsin gene from this library (data not shown). The number of the clones isolated confirmed that this BAC library covers almost 10x genome, since the *lws* gene was shown to be a single copy gene in cichlid from Lake Victoria (Terai et al., 2002c).

Recent studies from our laboratory (Takahashi et al., 2001a, 2001b, 2002; Terai et al., 2002a, 2002b) and others (Mayer et al., 1990; Sultmann et al., 1995; Nagl et al., 1998, 1999, 2000; Booton et al., 1999) have elucidated the close phylogenetic relationships among East African cichlid fishes. Remarkably, the genomes of species in Lake Victoria are nearly identical (Nagl et al., 1998, 1999; Terai et al., submitted). In fact, the DNA sequences of some housekeeping genes among several Haplochromine species are identical (Watanabe and Okada, unpublished data). Although nucleotide variations are present in a cichlid genome, most of them distribute as inter-specific polymorphisms among Haplochromine cichlids in Lake

Victoria. This is also an indication that the genomes of cichlids in Lake Victoria are essentially the same (Nagl et al., 1998, 1999; Terai et al., submitted). Thus, the BAC library constructed from *H. chilotes* may also prove to be useful for studies of the more than 200 Haplochromine cichlid species in Lake Victoria, although we must bear in mind that a very few genes responsible for speciation of *Haplochromis chilotes* are distinct from those of other species.

We thank Ms. M. Tochigi and Ms. H. Wada in RIKEN GSC for technical advices. This work was supported by a Grant-in-Aid for Scientific Research on Priority Areas (C) "Genome Science" from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

## REFERENCES

- Amemiya, C. T. Ota, T., and Lipman, G. W. (1996) In Nonmammalian genomic analysis: a practical Guide (eds. Birren, B. W. W. and Lai, E.), pp223–256, Academic Press, San Diego.
- Barel, C. D. N., Ligtvoet, W., Goldschmidt, T., Witte, F., and Goudswaard, P. C. (1991) The haplochromine cichlids in Lake Victoria: an assessment of biological and fisheries interests. In: Cichlid fishes: behaviour, ecology and evolution. (ed. Keenleyside, M. H. A.), pp. 259–279, Chapman and Hall, London.
- Booton, G. C., Kaufman, L., Chandler, M., Oguto-Ohwayo, R., Duan, W., and Fuerst, P. A. (1999) Evolution of the ribosomal RNA internal transcribed spacer one (ITS-1) in cichlid fishes of the Lake Victoria region. *Mol. Phylogenet. Evol.*

- 11, 273–282.
- Fryer, G. and Iles, T. D. (1972) The Cichlid Fishes of the Great Lakes of Africa. (eds.: Fryer G. and Iles, T. D.), TFH Publications, Neptune City, NJ.
- Fujiyama A, Watanabe H, Toyoda A, Taylor TD, Itoh T, Tsai SF, Park HS, Yaspo ML, Lehrach H, Chen Z, Fu G, Saitou N, Osoegawa K, de Jong PJ, Suto Y, Hattori M, Sakaki Y. (2002) Construction and analysis of a human-chimpanzee comparative clone map. *Science* **295**, 131–134.
- Greenwood, P. H. (1984) African cichlids and evolutionary theories. In: Evolution of fish species flock (eds.: Echelle and Kornfield), pp.141–154, University of Marine at Orono Press, Orono.
- Johnson, T. C., Scholz, C. A., Talbot, M. R., Kelts, K., Ricketts, R. D., Ngobi, G., Beuning, K., Ssemmanda, I. I., and McGill, J.W. (1996) Late Pleistocene Desiccation of Lake Victoria and Rapid Evolution of Cichlid Fishes. *Science* **273**, 1091–1093.
- Katagiri, T., Asakawa, S., Minagawa, S., Shimizu, N., Hirono, I., and Aoki, T. (2001) Construction and characterization of BAC libraries for three fish species; rainbow trout, carp and tilapia. *Anim. Genet.* **32**, 200–204.
- Majundar, K. C. and McAndrew, B. J. (1986) Relative DNA content of somatic nuclei and chromosomal studies in three genera, *Tilapia*, *Sarotherodon*, and *Oleochromis* of the tribe Tilapiini (Pisces, Cichlidae). **68**, 175–188.
- Mayer, A., Kocher, T. D., Basasibwaki, P. and Wilson, A. C. (1990) Monophyletic origin of Lake Victoria cichlid fishes suggested by mitochondrial DNA sequences. *Nature* **347**, 550–553.
- Nagl, S., Tichy, H., Mayer, W. E., Takahata, N. and Klein, J. (1998) Persistence of neutral polymorphisms in Lake Victoria cichlid fish. *Proc. Natl. Acad. Sci. USA* **95**, 14238–14243.
- Nagl, S., Mayer, W. E., and Klein, J. (1999) Isolation and sequencing of cDNA clones coding for the catalytic unit of glucose-6-phosphatase from two haplochromine cichlid fishes. *DNA Seq.* **10**, 25–29.
- Nagl, S., Tichy, H., Mayer, W. E., Takezaki, N., Takahata, N., and Klein, J. (2000) The origin and age of haplochromine fishes in Lake Victoria, east Africa., *Proc. R. Soc. Lond. B Biol. Sci.* **267**, 1049–1061.
- Shizuya, H., Birren, B., Kim, U. J., Mancino, V., Slepak, T., Tachiiri, Y., and Simon, M. (1992) Cloning and stable maintenance of 300-kilobase-pair fragments of human DNA in *Escherichia coli* using an F-factor-based vector. *Proc. Natl. Acad. Sci. USA* **89**, 8794–8797.
- Stiassny M. L. (1991) Phylogenetic intrarelationships of the family Cichlidae: an overview. In *Cichlid fishes: behaviour, ecology and evolution.* (ed. Keenleyside, M. H. A.), pp. 1–35, Chapman and Hall, London.
- Sultmann, H., Mayer, W. E., Figueroa, F., Tichy, H., and Klein, J. (1995) Phylogenetic analysis of cichlid fishes using nuclear DNA markers. *Mol. Biol. Evol.* **12**, 1033–1047.
- Takahashi, K., Nishida, M., Yuma, M., and Okada, N. (2001a) Retroposition of the AFC family of SINEs (short interspersed repetitive elements) before and during the adaptive radiation of cichlid fishes in Lake Malawi and related inferences about phylogeny. *J. Mol. Evol.* **53**, 496–507.
- Takahashi, K., Terai, Y., Nishida, M., and Okada, N. (2001b) Phylogenetic relationships and ancient incomplete lineage sorting among cichlid fishes in Lake Tanganyika as revealed by analysis of the insertion of retroposons. *Mol. Biol. Evol.* **11**, 2057–2066.
- Takahashi, K. and Okada, N. (2002) Mosaic structure and retropositional dynamics during evolution of subfamilies of short interspersed elements in African cichlids. *Mol. Biol. Evol.* **19**, 1302–1312.
- Terai, Y., Morikawa, N., Kawakami, K. and Okada, N. (2002a) Accelerated evolution of the surface amino acids in the WD-repeat domain encoded by the hagoromo gene in an explosively speciated lineage of East African cichlid fishes. *Mol. Biol. Evol.* **19**, 574–578.
- Terai, Y., Morikawa, N., and Okada, N. (2002b) The evolution of the pro-domain of bone morphogenetic protein 4 (bmp4) in an explosively speciated lineage of East African cichlid fishes. *Mol. Biol. Evol.* **19**, 1628–1632.
- Terai Y, Mayer WE, Klein J, Tichy H, Okada N. (2002c) The effect of selection on a long wavelength-sensitive (LWS) opsin gene of Lake Victoria cichlid fishes. *Proc Natl Acad Sci USA* **99**, 15501–15506.